

Balnase, a New Dimer-Forming Ribonuclease from *Bacillus altitudinis*

Dudkina E., Ulyanova V., Ilinskaya O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Current cancer treatments still remain ineffective and cause side effects which ruinously affect healthy cells. Among new promising anticancer drugs, special attention is paid to the ribonucleases (RNases) which possess selective cytotoxicity against malignant cells. It is found that besides enzymatic activity conformational state of RNase molecules plays a role in the anticancer activity. Balnase, a new RNase from *Bacillus altitudinis*, is a close homolog of *Bacillus pumilus* RNase (binase) which selectively kills malignant cells expressing oncogenes KIT, ras, AML1/ETO, and FLT3. Earlier, homogeneous sample of balnase was obtained. Here, we have characterized structural organization of balnase. It was shown that it is a natural dimer. The enzyme has the same conformational state as binase but balnase dimers are less stable.

<http://dx.doi.org/10.1007/s12668-016-0305-y>

Keywords

Anticancer activity, Balnase, Binase, Conformational state, Dimerization, Structural organization

References

- [1] Mitkevich, V., Ilinskaya, O., Makarov, A. (2015). Antitumor RNases: killer's secrets. *Cell Cycle*, 14, 931-932.
- [2] Mitkevich, V., Burnysheva, K., Ilinskaya, O., Pace, C., Makarov, A. (2014). Cytotoxicity of RNase Sa to the acute myeloid leukemia Kasumi-1 cells depends on the net charge. *Oncoscience*, 11, 738-744.
- [3] Ilinskaya, O., Dreyer, F., Mitkevich, V., Shaw, K., Pace, C., Makarov, A. (2002). Changing the net charge from negative to positive makes ribonuclease Sa cytotoxic. *Protein Science*, 11, 2522-2525.
- [4] Makarov, A., Kolchinsky, A., Ilinskaya, O. (2008). Binase and other microbial RNases as potential anticancer agents. *Bioessays*, 8, 781-790.
- [5] Ardelt, W., Ardelt, B., Darzynkiewicz, Z. (2009). Ribonucleases as potential modalities in anticancer therapy. *European Journal of Pharmacology*, 625, 181-189.
- [6] Zuo, Y., & Deutscher, M. (2002). Mechanism of action of RNase T. II. A structural and functional model of the enzyme. *The Journal of biological chemistry*, 277, 50160-50164.
- [7] Garvie, C., Vasanthavada, K., Xiang, Q. (2013). Mechanistic insights into RNase L through use of an MDMX-derived multi-functional protein domain. *Biochimica et Biophysica Acta*, 1834, 1562-1571.
- [8] Lin, R., Chien, H., Lin, S., Chang, B., Yu, H., Tang, W., et al. (2013). MCPIP-1 ribonuclease exhibits broad-spectrum antiviral effects through viral RNA binding and degradation. *Nucleic Acids Research*, 41, 3314-3326.
- [9] Gotte G., Mahmoud Helmy A., Ercole C., Spadaccini R., Laurents D., Donadelli M., Picone D. (2012) Double domain swapping in bovine seminal RNase: formation of distinct N- and C-swapped tetramers and multimers with increasing biological activities. *PloS One* 7: doi:10.1371/journal.pone.0046804

- [10] Ilinskaya, O., Decker, K., Koschinski, A., Dreyer, F., Repp, H. (2001). Changing the net charge from negative to positive makes ribonuclease Sa cytotoxic. *Toxicology*, 156, 101-107.
- [11] Mitkevich, V., Petrushanko, I., Kretova, O., Zelenikhin, P., Prassolov, V., Tchurikov, N., et al. (2010). Oncogenic c-kit transcript is a target for binase. *Cell Cycle*, 9, 2674-2678.
- [12] Mitkevich, V., Petrushanko, I., Spirin, P., Fedorova, T., Kretova, O., Tchurikov, N., et al. (2011). Sensitivity of acute myeloid leukemia Kasumi-1 cells to binase toxic action depends on the expression of KIT and AML1-ETO oncogenes. *Cell Cycle*, 10, 4090-4097.
- [13] Mitkevich, V., Schulga, A., Trofimov, A., Dorovatovskii, P., Goncharuk, D., et al. (2013). Structure and functional studies of the ribonuclease binase Glu43Ala/Phe81Ala mutant. *Acta Cryst D Biol Crystallogr*, 69, 991-996.
- [14] Dudkina, E., Kayumov, A., Ulyanova, V., Ilinskaya, O. (2014). New insight into secreted ribonuclease structure: binase is a natural dimer. *PloS One*, 12, e115818.
- [15] Dementiev, A., Orlov, V., Shlyapnikov, S. (1993). Complete primary structure of an extracellular ribonuclease of *Bacillus thuringiensis*. *Russ J Bioorganic Chem*, 19, 853-861.
- [16] Dudkina, E., Ulyanova, V., Shah, M. R., Khodzhaeva, V., Dao, L., Vershinina, V., et al. (2016). Three-step procedure for preparation of pure *Bacillus altitudinis* ribonuclease. *FEBS Open Bio*, 6, 24-32.
- [17] Ilinskaya, O., Karamova, N., Ivanchenko, O., Kipenskaya, L. (1996). SOS-inducing ability of native and mutant microbial ribonucleases. *Mutation Research*, 354, 203-209.
- [18] Laemmli, U. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.